

RELAXATION DIELECTRIC PROPERTIES OF GPI CRYSTAL

I.R. Zacheck, O.B.Bilenka

Lviv Polytechnic National University, Lviv,

Within pseudospin model of ferroelectric GPI on the basis of Zubarev nonequilibrium statistical operator [1] we obtained kinetic equation for pseudospin operators [2]. Solving this equation we have calculated the temperature and frequency dependences of dynamic dielectric permittivity (fig.1, 2) of the crystal, relaxation time. At the proper set of parameters satisfactory description of experimental data is obtained.




Figure 1. Temperature dependences of ϵ'_{33} and ϵ''_{33} for various frequencies v (MHz):

1.0 – 1, \diamond [3]; 15.0 – 2, \circ [3]; 230.0 – 3, \square [3]; 610 – 4, \square [3]; 2000 – 5, \square [3]; 27000 – 6, \triangleright [3].




Figure 1. Frequency dependences of ϵ'_{33} and ϵ''_{33} of GPI crystal for various ΔT (K):

1.0 – 1, 2.0 – 2, 5.0 – 3, 10 – 4; Symbols: \bullet [4], \blacksquare [3], \blacklozenge [5], \blacktriangleright [6], \blacktriangle [7].

2. [1] D.N. Zubarev. Nonequilibrium statistical thermodynamics. –M.: Nauka, 1971. -416p.
3. [2] G.O. Berim, A.R. Kessel. Physica. 101A, N 1, 1980, p. 112.
4. [3] R. Tchukvinskyi, Z. Czapla, R. Sobiestianskas, A. Brilingas, J. Grigas, J. Baran, Acta Phys. Polonica A, 92, 1997, p. 1191.
5. [4] M. Wiesner, Phys.stat. sol (b), 238, 2003, p. 68.
6. [5] J. Baran, G. Bator, R. Jakubas, M. Sledz, J. Phys.: Condens Matter, 8, 1996, p.10647.
7. [6] J. Nayeem, H. Wakabayashi, T. Kikuta, T. Yamazaki, N. Nakatani, Ferroelectrics, 269, 2002, p.153.
8. [7] S. Dacko, Z. Czapla, J. Baran, M.Drozdz, Physics Letters A, 221, 1996, p.217.